



Spontaneous Coronary Artery Dissection: An Elusive Disease of the Coronaries

SMITHA NARAYANA GOWDA, MD 

SWAMINATHAN PERINKULAM SATHYANARAYANAN, MD 

ALPESH SHAH, MD 

*Author affiliations can be found in the back matter of this article

POINTS TO REMEMBER



The articles in our Points to Remember column highlight important “need to know” facts about conditions that cardiovascular healthcare professionals may encounter. These may come from any medical specialty, such as nephrology or neurology. The article in this issue is provided by Swaminathan Perinkulam Sathyanarayanan, MD, Department of Internal Medicine, University of South Dakota, and Smitha Narayana Gowda, MD, and Alpesh Shah, MD, Houston Methodist DeBakey Heart & Vascular Center, Houston Methodist.

CORRESPONDING AUTHOR:

Alpesh Shah, MD

Houston Methodist DeBakey Heart & Vascular Center, Houston Methodist Hospital, Houston, Texas, US

ARShah@houstonmethodist.org

KEYWORDS:

spontaneous coronary artery dissection; myocardial infarction; acute coronary syndrome

TO CITE THIS ARTICLE:

Gowda SN, Sathyanarayanan SP, Shah A. Spontaneous Coronary Artery Dissection: An Elusive Disease of the Coronaries. *Methodist DeBakey Cardiovasc J.* 2022;18(3):89–93. doi: [10.14797/mdcvj.1129](https://doi.org/10.14797/mdcvj.1129)




Figure 3 (A) Spontaneous coronary artery dissection (SCAD) of ramus coronary artery. **(B)** Possible SCAD in the apical branch of bifid left anterior descending coronary artery (LAD) and healing of previous SCAD in diagonal branch of bifid LAD.

- Between 1% and 4% of ACS cases are from SCAD. An important cause of myocardial infarction (MI) in young women, it accounts for up to 35% of ACS cases in women < 50 years of age and is the most common cause of pregnancy-associated MI.¹
- Risk factors include female gender, pregnancy, fibromuscular dysplasia connective tissue disorders, hormonal therapies, corticosteroids, and emotional and physical stressors. Traditional risk factors of MI are generally few or absent.³
- Chest pain, the most common presentation of SCAD, rarely presents with cardiogenic shock, ventricular arrhythmias, and sudden cardiac death.²
- Diagnosis of SCAD is generally made on coronary angiography. Based on its angiographic appearance (Figure 4), it has three types of Yip-Saw classification: (1) multiple radiolucent lumens or arterial wall contrast staining; (2) presence of diffuse stenosis of varying severity and length (most commonly seen); and (3) focal or tubular stenosis mimics atherosclerosis.⁴
- During diagnostic uncertainty, intracoronary imaging modalities such as optical coherence tomography (OCT) and intravascular ultrasound (IVUS) should be considered after weighing risk versus benefits because these maneuvers can cause extension of dissection, occlusion of the lumen, and iatrogenic dissection.²
- While optimal management is up for debate, conservative medical therapy is recommended in most cases. Healing of SCAD lesion occurs in a majority of patients (70–97%) after conservative management, but recurrence is common.¹
- Revascularization with percutaneous coronary intervention or coronary artery bypass graft surgery should be considered in patients with symptoms of ongoing ischemia, hemodynamic instability, or left main dissection and/or involvement of the proximal part of the epicardial vessels.¹
- Percutaneous coronary intervention (PCI) may be technically challenging due to difficulty assessing the true lumen and propagation of dissection antegrade and retrograde. PCI should be undertaken at a high experience center with IVUS or OCT guidance and with careful attention to complete coverage of the entire length of the lesion, especially the proximal origin of dissection.
- Long-term medical therapy includes aspirin and beta blockers. Beta blocker use has been associated with lower risk of recurrent SCAD. The role of antiplatelets, like clopidogrel, remains unclear but is used commonly in contemporary clinical practice.⁵

COMPETING INTERESTS

The authors have no competing interests to declare.

